



Editorial

Copyright © All rights are reserved by Diego Liberati

How to Measure Our Sensitivity in Order to Better Understand What Happens and Decide What to Do

Diego Liberati*

Consiglio Nazionale delle Ricerche, Italy

*Corresponding author: Diego Liberati, Consiglio Nazionale delle Ricerche, Italy.

Received Date: August 12, 2018

Published Date: August 28, 2018

Editorial

We are obviously sensitive to every kind of stressors. To measure effects is of paramount necessity. A first not specific but quite sensitive way to detect and quantify stress is heart rate variability: sympatho-vagal balance is known to be altered either in acute or even chronically according to the state and of the dynamics of each of us; specificity could be at least partially achieved by keeping fixed as much as possible every other stimuli [1]. Less immediate, and still originally a-specific, is the analysis of the central nervous system instead of the autonomous one: coherences among brain areas, investigated via EEG, MEG, fMRI, NIRS, do even account on our plasticity to the change [2]. Deconvolution of blood samples may help in noninvasively assessing un-accessible and nano-metric pituitary secretion in controlling hormone loops [3]. A real precision analysis is needed to achieve very specific results: epigenetics makes us enhancing gene mediated protein expression in such a way that salient involved genes are detectable in assays together with their networking behavior with proteins expressed to face stimuli [4]. Modeling biophysical and biochemical interactions at molecular, domain and even atomic scale could become the ultimate level in approaching the effect from macro to meso, coarse and micro scales [5,6].

References

1. Pagani M, Mazzuero G, Ferrari A, Liberati D, Cerutti S et al. (1991) Sympathovagal interaction during mental stress: a study employing spectral analysis of heart rate variability in healthy controls and in patients with a prior myocardial infarction. *Circulation* 83(4) II: 43-51.
2. Baraldi P, Manginelli AA, Maieron M, Liberati D, Porro CA (2007) An ARX model-based approach to trial by trial identification of fMRI-BOLD responses. *NeuroImage* 37(1): 189-201.
3. De Nicolao G, Liberati D (1993) Linear and Nonlinear Techniques for the Deconvolution of Hormone Time-Series. *IEEE Transactions on Biomedical Engineering* 40 (5): 440-455.
4. Muselli M, Liberati D (2002) Binary rule generation via Hamming Clustering. *IEEE Transactions on Knowledge and Data Engineering* 14(6): 1258-1268.
5. Mezza D, Pappalettera M, Liberati D (2016) A quantitative numerical model for TNF- α mediated cellular Apoptosis. *Biol Eng Med* 1(1): 1-8.
6. Sacco E, Farina M, Greco C, Lamperti S, Busti S, et al. (2012) Regulation of hSos1 activity is a system-level property generated by its multi-domain structure. *Biotechnol Adv* 30(1): 154-168.