



Research Article

Copyright © All rights are reserved by Elizaveta I Bon

Violations of the Ultrastructure of Neurons in the Cerebral Cortex of Rats with Total Cerebral Ischemia

Elizaveta I Bon*, Maksimovich N Ye, Zimatkin SM, Ostrovskaya OB, Smirnov V Yu, Nosovich MA and Khrapovitskaya KA

Grodno State Medical University, Grodno, Republic of Belarus

Corresponding author: Elizaveta I Bon, Grodno State Medical University, Grodno, Republic of Belarus.

Received Date: December 20, 2022

Published Date: January 09, 2023

Annotation

The ultrastructural characteristics of neuronal organelles are important indicators of the degree of brain damage during ischemic exposure, reflecting the severity of compensation, which necessitates the study of changes in the ultrastructure of brain neurons. With total cerebral ischemia, organelles are disorganized both in the neurons of the parietal cortex and in the neurons of the hippocampus, causing profound disturbances in energy production and metabolism.

Keywords: Neurons; Parietal cortex; Hippocampus; Ischemia

Introduction

With cerebral ischemia, a chain of pathogenetic disorders develops in its structures, among which one of the leading ones is energy deficiency, which leads to the development of cellular pathology due to violations of homeostasis, enzyme activity, membrane integrity and energy pumps. Under conditions of cerebral ischemia, synaptic transmission mechanisms are selectively disrupted, which contributes to disruption of autoregulation of local blood flow, the development of vasospasm, increased platelet aggregation and the development of intravascular stasis, aggravating hypoxia and increasing energy deficiency [1-5]. The ultrastructural characteristics of neuronal organelles are important indicators of the degree of brain damage during ischemic exposure, reflecting the severity of compensation, which necessitates the study of changes in the ultrastructure of brain neurons. Currently, there are no data on the severity of these disorders depending on the type of ischemic injury and its severity. These studies are

relevant, as they allow us to study the nature of disorders of brain neurons at the ultrastructural level, depending on the severity of ischemia.

Materials and Methods of Research

The experiments were carried out on 12 male outbred white rats weighing 260 ± 20 g in compliance with the requirements of the Directive of the European Parliament and of the Council No. 2010/63/EU of September 22, 2010 on the protection of animals used for scientific purposes. Cerebral ischemia modeling was carried out under conditions of intravenous thiopental anesthesia (40-50 mg/kg). The studies used models of total cerebral ischemia. The control group included 4 males, the ischemia group included 8 males.

Total cerebral ischemia was modeled by decapitation of animals. The brain sampling was carried out 1 hour and 24 hours

after decapitation. The control group consisted of sham-operated rats of similar sex and weight. Electron microscopic studies were performed in the parietal cortex and hippocampus of the brain of rats.

Immediately after decapitation and quick extraction of the brain, sections were cut out with a blade and placed in 1% osmium fixative in Millonig's buffer (pH=7.4) for 2 hours at a temperature of 4°C. Next, the sections were washed in a mixture of Millonig's buffer (20 ml) and sucrose (900 mg), dehydrated in alcohols of increasing concentration, a mixture of alcohol and acetone, pure acetone; passed through a mixture of resins (araldite M + araldite H + dibutyl phthalate + DMR-30) and acetone and embedded in a mixture of resins. Semi-thin sections (about 350 nm thick) were made on an MT-7000 ultramicrotome (RMC, USA), stained with methylene blue, and the sections necessary for study were cut out with a blade.

Ultrathin sections (about 35 nm thick) were made on the same ultramicrotome, assembled on support grids, and counterstained with uranium acetate and lead citrate. To do this, the meshes with sections were dipped into a drop of uranyl acetate and kept for 20 minutes in the dark at room temperature, then washed in 3 portions of bidistilled water for 5 seconds and contrasted with lead citrate for 8 minutes, washed in 3 portions of bidistilled water for 5 seconds.

The resulting preparations were studied under a JEM-1011

electron microscope (JEOL, Japan) and photographed with an Olympus MegaView III digital camera (Olympus Soft Imaging Solutions, Germany).

Morphometry of ultrastructures was performed using the Image Warp program (Bit Flow, USA). Density, size, shape of organelles, density and length of mitochondrial cristae, sizes of lysosomes and their density, the number of endoplasmic reticulum-bound and free ribosomes were measured, and their ratio was calculated.

To prevent a systematic measurement error, brain samples from the compared control and experimental groups of animals were studied under the same conditions. As a result of the research, quantitative continuous data were obtained. Since the experiment used small samples that had a non-normal distribution, the analysis was performed by nonparametric statistics using the licensed computer program Statistica 10.0 for Windows (StatSoft, Inc., USA). The data are presented as Me (LQ; UQ), where Me is the median, LQ is the value of the lower quartile; UQ is the value of the upper quartile. Differences between groups were considered significant at $p < 0.05$ (Kruskell-Wallis test with Bonferoni correction).

Research Results

The characteristics of changes in the ultrastructure of neurons in the cerebral cortex of rats with total cerebral ischemia are shown in (Table 1).

Table 1: Parameters of ultramicroscopic morphometry of neuron organelles in the parietal cortex and hippocampus of rats with total cerebral ischemia (TCI), Me(LQ;UQ).

Index		Parietal cortex			Hippocampus			
		Control	TCI 1 hour	TCI 1 day	Control	TCI 1 hour	TCI 1 day	
Mito-	density		1,8(1,7;2,2)	1,6(1,5;1,7)	1,2(1,0;1,3) **	2,1(1,7;2,2)	1,7(1,6;1,7)	1,3(1,1;1,4) **
	area, μm^2		0,26 (0,17;0,37)	0,11(0,06;0,12) *	0,10(0,08;0,12) *	0,21(0,17;0,26)	0,10(0,08;0,12) *	0,09(0,08;0,11) *
chondria	Form factor, unit		0,63(0,61;0,72)	0,87(0,78;0,96)*	0,93(0,83;0,96)	0,71(0,60;0,75)*	0,86(0,84;0,95)*	0,89(0,85;0,93)*
	Elongation factor, units		3,8(3,5;4,1)	1,4(1,3;1,5)	1,4(1,3;1,5)	2,1(1,9;2,5)	1,5(1,3;1,7)*	1,4(1,2;1,5)*
	mitochondrial crist density		76(71;82)	45(41;51)*	40(37;45)*	62(59;72)	45(41;52)*	40(37;45)*
	length of mitochondrial cristae / μm^2		12(10;15)	5(4;6)*	4(3;6)*	13(12;18)	6(4;7)*	5(4;7)*
Ribo-	amount / μm^2		20,9(19,3;22,7)	17,6(16,5;18,7) *	17,4(16,8;19,2)	20(18,1;22,8)	16,2(14,6;19,4) *	16,3(15,0;18,7)*
	free / μm^2		4,7(4,1;5,8)	16,0(15,3;16,8)*	16,5(16,0;17,8)*	6,0(4,8;7,3)	14,5(13,3;17,3)*	15,5(14,3;17,5)*
	bound / μm^2		16,2(15,2;16,9)	1,6(1,2;1,9)*	0,9(0,8;1,4)*	14,0(13,3;15,5)	1,7(1,3;2,1)*	0,8(0,7;1,2)*
somes	ratio of bound and free ribosomes		3,4	0,1*	0,05*	2,3	0,1*	0,05*
Lyso-	density		0,4(0,3;0,5)	0(0;0,1)	0(0;0)	0,5(0,4;0,6)	0(0;0,1)	0(0;0)
	area, μm^2		0,02(0,01;0,03)	-	-	0,03(0,02;0,04)	-	-

Note* - $p < 0.05$ compared with the control table, + - compared with the "TCI 1 hour" group.


When studying the ultrastructure of neurons, there was no change in the number of mitochondria in the cytoplasm of rat neurons during 1-hour cerebral ischemia ($p > 0.05$), but there

was a change in their size and shape. With 1-hour TCI, compared with the indicators in the control group, in the parietal cortex, the mitochondrial area decreased by 58% ($p < 0.05$), the elongation

factor (organelle elongation) decreased by 63% ($p < 0.05$), and the form factor (roundness of the organelle) increased by 28%, ($p < 0.05$). Similar changes were noted in the hippocampus: a decrease in the area and elongation factor by 52% ($p < 0.05$) and by 29% ($p < 0.05$), respectively, and an increase in the form factor by 14% ($p < 0.05$), which indicates rounding of mitochondria in both studied structures.


The density of mitochondrial cristae and their length decreased in the parietal cortex by 40% ($p < 0.05$) and 58% ($p < 0.05$), respectively. In the hippocampus, these indicators decreased by 27% ($p < 0.05$) and 54% ($p < 0.05$), respectively. With TCI with an ischemic period of 1 day, there was a decrease in mitochondrial density: by 33% in the parietal cortex ($p < 0.05$), and by 38% in the

hippocampus ($p < 0.05$), compared with the control group and by 25%, ($p < 0.05$), and 24%, ($p < 0.05$), respectively, compared with the "TCI 1 hour" group. At the same time, in the parietal cortex, the area of mitochondria decreased by 59% ($p < 0.05$), the elongation factor decreased by 63% ($p < 0.05$), and the form factor, on the contrary, increased by 32% ($p < 0.05$). In the hippocampus, the area decreased by 53% ($p < 0.05$), the elongation factor decreased by 33% ($p < 0.05$), while the form factor increased by 20% ($p < 0.05$). The density of mitochondrial cristae and their length were less in the parietal cortex by 47% ($p < 0.05$) and 67% ($p < 0.05$), respectively. In the hippocampus, these indicators decreased by 35% ($p < 0.05$) and 62% ($p < 0.05$), respectively. There were no differences in these parameters between the groups "TCI 1 hour" and "TCI 1 day" ($p > 0.05$) (Figures 1,2).



Electrongram. Magnification 50000

Figure 1: Mitochondria of neurons in the parietal cortex of the rats of the control group.



Electrongram. Magnification 50000


Figure 2: Mitochondria of neurons in the parietal cortex of rats with 1-hour total cerebral ischemia.

Compared with the indicators in the "control" group, the number of free ribosomes during 1-hour TCI increased by 71%

in the parietal cortex ($p < 0.05$), and in the hippocampus - by 57% ($p < 0.05$). The ratio of the number of bound and free ribosomes


decreased from 3.43 in the control group to 0.11 in the parietal cortex ($p < 0.05$) and from 2.33 to 0.12 in the hippocampus ($p < 0.05$). With 1-day TCI, the number of free ribosomes in the cytoplasm of neurons of the parietal cortex and hippocampus increased by 72% ($p < 0.05$) and by 58% ($p < 0.05$) compared to the values in the control group, which did not differ from changes in TCI lasting 1 hour ($p > 0.05$). The ratio of the number of bound and free ribosomes decreased from 3.4 in the control group to 0.05 in the parietal cortex ($p < 0.05$) and from 2.33 to 0.05 in the hippocampus ($p < 0.05$).

There was a significant decrease in the density of lysosomes. With 1-hour TCI, only single lysosomes were observed in the cytoplasm of neurons ($p < 0.05$). With 1-day TCI, lysosomes were absent in both studied parts of the brain ($p < 0.05$). Along with this, compared with the control group, there was a change in the expansion and vacuolization of the cisterns of the granular endoplasmic reticulum in TCI with a predominance of free ribosomes in the cytoplasm, (Figures 3,4).




Electrongram. Magnification 50000

Figure 3: Endoplasmic reticulum of neurons in the parietal cortex of the control group rats.




Electrongram. Magnification 50000

Figure 4: Endoplasmic network of neurons in the parietal cortex of rats with 1-hour total cerebral ischemia.



Electronogram. Magnification 50000

Figure 5: Golgi complex of neurons in the pyramidal layer of field CA1 of the hippocampus of rats in the control group.



Electronogram. Magnification 50000

Figure 6: Golgi complex of neurons in the pyramidal layer of the field CA1 of the hippocampus of rats with 1-day total cerebral ischemia.

Deformation and vacuolization of the Golgi complex was also noted in comparison with the control group, (Figures 5,6).

Ultramicroscopic morphometry of organelles revealed no differences between the groups "TCI 1 hour" and "TCI 1 day" ($p > 0.05$), except for a decrease in the density of mitochondria with 1-day TCI, which indicates a progressive disorder of these organelles, leading to an increase in energy shortages. Discussion of the obtained data. According to the literature data, changes in the mitochondria of neurons are noted during cerebral hypoxia: swelling, destruction of their cristae, uneven distribution in the cytoplasm, which indicates a violation of the energy supply of neurons [1,2,3,4,6]. The destruction of the outer and inner membranes leads to a decrease in the number of mitochondria due to a violation of the permeability for cations, which leads to edema and rupture. Active swelling of mitochondria is associated with the operation of the electron transport chain. There is an

expansion of the tubules of the granular and smooth endoplasmic reticulum, a change in their structure, disintegration into small granules, the appearance of large vacuoles and loops [5,6,7,8]. Free ribosomes predominate, forming extensive clusters. This is one of the manifestations of the energy deficiency that is formed in the cell, since the fixation of ribosomes to the membranes of the rough endoplasmic reticulum with the participation of the ribophorin protein is an energy-dependent process. Under conditions of ischemic exposure, the neuron reduces protein export and seeks to direct it for internal needs. Disorganization of the granular endoplasmic reticulum leads to the accumulation of newly formed proteins in the cytoplasm. As cellular hypoxia and acidosis increase, protein denaturation increases [9]. Due to the accumulation of water, the expansion of the cisterns of the Golgi complex and their partial fragmentation are noted [7,10]. The average number of lysosomes increases, their sizes increase. There is an exit into the cytoplasm

and activation of hydrolytic enzymes of lysosomes - cathepsin, ribonuclease, acid phosphatase, deoxyribonuclease, hyaluronidase and other enzymes, which leads to autolysis [4,5,8,9,11].

The data obtained correspond to the literature data, supplementing them with the features of the development of organelle disorganization during ischemia of varying severity.

Conclusion

Thus, with total cerebral ischemia, organelles are disorganized both in the neurons of the parietal cortex and in the neurons of the hippocampus, causing profound disturbances in energy production and metabolism.

Funding source

This study was not supported by any external sources of funding. Competing interests. The authors declare that they have no competing interests.

Author contribution

All authors confirm the compliance of their authorship, according to international ICMJE criteria (all authors made a substantial contribution to the conception of the work, acquisition, analysis, interpretation of data for the work, drafting and revising the work, final approval of the version to be published).

References

1. Hong A, Bing Zh, Xunming J (2021) Mitochondrial quality control in acute ischemic stroke. *J Cereb Blood Flow Metab.* 41(12): 3157-3170.
2. Nicholls DG, Budd SL (2000) Mitochondria and neuronal survival. *Physiol Rev* 80(1): 315-360.
3. Bon' EI, Zimatkin SM (2017) Dark brain neurons. *Morphology* 6: 81-86.
4. Nina J Solenski, Charles G diPierro, Patricia A Trimmer, Aij-Li Kwan, Gregory A Helms (2002) Ultrastructural Changes of Neuronal Mitochondria After Transient and Permanent Cerebral Ischemia. *Stroke* 33(3): 816-824.
5. Mattson MP, Clumsee C, Yu ZF (2000) Apoptotic and antiapoptotic mechanisms in stroke. *Cell Tissue Re* 301: 173-187.
6. Zhi H, Niya N, Qiongxiu Zh, Seyed Esmaeil Kh, Maryam Farzaneh (2020) Mitochondria as a therapeutic target for ischemic stroke. *Free Radic Biol Med* 146: 45-58.
7. Garcia JH, Lossinsky AS, Kauffman FC, Conger KA (1978) Neuronal ischemic injury: light microscopy, ultrastructure and biochemistry. *Acta Neuropathologica* 43: 15-24.
8. Snider BJ, Gottron FJ, Choi DW (2004) Apoptosis and necrosis in cerebrovascular disease. *Ann NY Acad Sci* 893: 243-253.
9. Colbourne F, Sutherland GR, Auer RN (1999) Electron microscopic evidence against apoptosis as the mechanism of neuronal death in global ischemia. *J Neurosci* 19: 4200-4210.
10. Belenichev IF (2014) Neuroprotection and neuroplasticity: monograph. K: Polygraph Plus LLC: 512.
11. Zimatkin SM (2017) Maternal alcohol intake induces dramatic ultrastructural changes in offspring brain cortex neurons. *Brain and Nerves* 1(1): 1-4