

Ring Models of Atoms and Molecules with Two Opposite Spins of Proton and Neutron Rings (Atom Type A and Atom Type B)

Pavel Ošmera*

Mechanical Department, Brno University of Technology, Czech Republic

*Corresponding author: Pavel Ošmera, Mechanical Department, Brno University of Technology, Czech Republic.

Received Date: April 20, 2023

Published Date: May 18, 2023

Abstract

Using the ring model of atoms and molecules, it was revealed that there must be two types (type A and type B) of atoms that differ in the direction of rotation of individual rings (for type B, all rings rotate opposite to the directions of rotation for type A). The rotations of protons and neutrons in atoms are given by a system that resembles a "cogwheel" in gear box.

Keywords: Ring model; Direction of rotation of proton; Neutron and electron; Directions of magnetic moments; Photocatalytic cleaner with TiO_2

Introduction

The solution to this problem was found in the design of the TiO_2 model, which has a total of 22 protons, 22 electrons and 26 neutrons in titanium atom (Ti). Each oxygen atom has 8 protons, 8 electrons and 8 neutrons. The TiO_2 model therefore consists of 118 rings (protons, neutrons and electrons). Two types of electrons are distinguished in the model (valence electrons are colored light blue and inner electrons are colored dark blue). The direction of rotation of the proton determines the direction of the magnetic moment that exits the atom. For an external view of the proton, it has (for clockwise rotation of the proton) the proton has a magnetic moment that exits the proton (yellow arrow). For counter clockwise rotation of the proton) the proton has a magnetic moment that

enters the proton (green arrow). The article uses the knowledge described in [1-8].

Model of the A-Type and B-Type Oxygen Atom

In Figure 1 is a topological model of oxygen of type A. The topological model follows the mutual connections of the rings in the atom and molecule, while not observing the correct dimensions and distances of the individual rings. For example: at the drawn size of the proton, the electron would be far away on the model (it would not fit in the A4 format). On the contrary, with the correct size of the distance between the electron and the proton, the proton would be a dot in the picture (the nucleus of the atom would also be a dot) (Figure 1 and Figure 2).




Figure 1: Topological model of an A-type oxygen atom (red rings are protons, yellow are neutrons, blue are electrons, yellow and green arrows are directions of magnetic moments).




Figure 2: Topological model of the B-type oxygen atom (red rings are protons, yellow are neutrons, blue are electrons, yellow and green arrows are directions of magnetic moments).

Oxygen Molecule O_2




Figure 3: Topological model of the oxygen molecule: connection of type A and type B atoms (red rings are protons, yellow are neutrons, blue are electrons, yellow and green arrows are directions of magnetic moments)

In the case of an oxygen molecule, oxygen atoms of type A and type B are joined. The magnetic moments in the binding of two protons must follow the same direction. Only the magnetic moment with the yellow arrow can be combined with the magnetic moment with the green arrow. The hydrogen molecule is in Figure 3.

Name of Electrons Inside Atom of Oxygen

Figure 4 Name of electrons inside two type A and B atoms of oxygen.




Figure 4: Name of electrons inside two type A and B atoms of oxygen.

Type A and type B 3 of carbon atoms

Figure 5 shows both versions of the carbon atom (type A and type B). Since the carbon atom is centripetal, we also get type B from type A by rotating the type A model by 180 degrees (Figure 5).

Carbon dioxide CO₂ molecule

Figure 6 shows a model of the carbon dioxide molecule, which consists of a carbon atom and two oxygen atoms of both types.

Carbon dioxide has both types of oxygen atom in the molecule. Two oxygen atoms of the same type cannot form a carbon dioxide molecule. Molecule CO₂ has linear structure (Figure 6).

Titanium dioxide TiO₂ molecule

Figure 7 shows a model of titanium atom and the titanium dioxide molecule, which consists of a titanium atom and two oxygen atoms of both types. Titanium dioxide has both types of

oxygen atom in the molecule. Two oxygen atoms of the same type cannot form a carbon dioxide molecule. TiO_2 molecule has linear structure as CO_2 molecule. TiO_2 molecule is used in photocatalytic

cleaner. The surface with TiO_2 is activated using UV light (band A – 365 nm) emitted by UV LEDs (Figure 7).




Figure 5: Topological model of type A and type B carbon atoms (red rings are protons, yellow are neutrons, blue are electrons, yellow and green arrows are directions of magnetic moments).




Figure 6: Topological linear model of the carbon dioxide molecule (red rings are protons, yellow are neutrons, blue are electrons, yellow and green arrows are directions of magnetic moments).




Figure 7: Topological model of titanium atom and linear model the titanium dioxide molecule (red rings are protons, yellow are neutrons, blue are electrons, yellow and green arrows are directions of magnetic moments).

Photocatalytic Cleaner with Titanium Dioxide




Figure 8: Photocatalytic cleaner 2 HELICO.

In Figure 8 is a look inside the photocatalytic air cleaner 2HELICO, manufactured by NCT s.r.o. (New Czech Tech. Ltd.). Using the dog cleaner (purifier) breaks down complex molecules (viruses, bacteria, fungi, cat and dog odors, cigarette smoke) into harmless substructures (Figure 8).

Conclusion

The aim of the article was to show the necessity that there must exist for each atom both types (type A and type B) in which all the rings rotate in the opposite direction. Therefore, protons have magnetic moments with opposite directions, which enables them to combine into molecules. Only protons that have opposite directions of magnetic moments can be connected (only connecting the direction with the green arrow to the direction with the yellow arrow). We can call this direction: spin in and spin out. Knowledge of structure rules of the atomic nucleus and the properties of vortex electromagnetic field allow us to create relatively precisely the structures of individual atoms and molecules. Properties of atoms are largely described by the structure of their electron shells. However, the standard model of atoms does not allow define this structure exactly. New theory VFRT (vortex-fractal-ring-theory) can solve this lack. Theory VFRT uses fractal ring structure of the electron, the proton and the neutron, and can describe the inner structure of atomic nuclei. Fractal descriptions of Nature are very promising. The atomic nucleus can be built from the ring protons and neutrons. This new theory assumes that the arrangement of electron shells arises from the structure of the atomic nucleus. Electrons are not in orbit around the atomic nucleus, but each electron levitates with the corresponding proton of the nucleus. The levitation bond between the electron and the proton is formed by an electromagnetic vortex structure. Theory VFRT expands understanding of nature through a new perspective on the evolution

of lifeless nature using a vortex, fractal ring substructures with self-organization, from quarks, electrons, protons and neutrons, atoms, molecules, to the structure of complex organic compounds.

Acknowledgement

None.

Conflict of Interest

No Conflict of interest.

References

1. Pauling L (1988) General Chemistry, Dover publication, Inc, New York, USA.
2. Ramsden EN (2000) A-Level Chemistry, Nelson Thornes Ltd, fourth edition, USA.
3. Ošmera P (2006) Chaotic system with vortex-fractal structures, in Proceedings of 13th Zittau East-West Fuzzy Colloquium.
4. Ošmera P (2012) Fractal dimension of electron, published in the Proceedings of MENDEL, page 186-191.
5. Ošmera P (2015) Ring structures of atoms and molecules, 2015-08-9, San Diego, California, USA.
6. Ošmera P (2016) RING STRUCTURES OF MATTER, 2017-10-18, published in the Proceedings of NANOCON2017, Brno, Czech Republic, USA.
7. Ošmera P, Werner P (2015) Ring structure of atoms and molecules, Proceedings of SPIE Vol.95701C-1, San Diego, USA.
8. Werner P (2018) Fundamentals of modeling the ring structure of elementary particles of matter, Institute of Theoretical and Experimental Electrical Engineering, Brno, USA.
9. More about VFRT theory can be found on <http://www.pavelosmera.cz>