

**Short Communication**

Copyright © All rights are reserved by Castaldi M

# Oral health and physical activity

**Castaldi M<sup>1\*</sup> and Nardi GM<sup>2</sup>**<sup>1</sup>RDH DHA, ATASIO Active Member<sup>2</sup>RDH DHA, Associate Professor, Department of Oral and Maxillofacial Sciences, Sapienza University of Rome, Italy**\*Corresponding author:** Castaldi Matteo, Public Health and Epidemiology Department, academy of advanced technologies in oral hygiene sciences, Italy.**Received Date: February 09, 2023****Published Date: March 07, 2023**

## Introduction

Oral health is an important element of overall health, well-being, and quality of life. The nexus between sport and oral health has largely been investigated through studies focused on the risk of trauma, dental caries, dental erosion, and periodontal inflammation [1]. Poor oral health can reduce quality of life and induce a systemic inflammatory response [2-5]. Thus, poor oral health could affect athletic performance. However, the relationship between oral health and performance is not well understood [6-8].

Numerous studies report a relationship between strenuous physical activity and problems referred to the oral cavity. One of these systematic reviews [9] describes that dental caries, periodontal disease, dental erosion and pericoronitis (infections around wisdom teeth)/impacted third molars were widely represented in the oral cavity of athletes. The range of proportion of athletes affected by these conditions were dental caries 15-75%, moderate-to-severe periodontitis up to 15%, dental erosion 36-85% and pericoronitis/impacted third molars 5-39%. Dental trauma was reported by 14-57% of athletes in at-risk sports. Disease incidence was generally not clearly differentiated by socioeconomic status; however, poor oral health appears to affect athletes both from developing and developed countries.

The available evidence suggests that poor oral health is common in elite or professional athletes and the oral health of elite athletes is like non-athlete disadvantaged populations [9].

## Performance

Several studies have investigated whether oral health has measurable impacts on sport performance. Realistically, this

is a difficult aspect to measure. Whilst dental and oro-facial trauma or severe infection can take an athlete out of training or competition (time loss injury) other health impacts (including from oral health) are likely to be of less magnitude and therefore challenging to measure. Although smaller in magnitude, these impacts could be important to training and performance and could be considered like the potential effect of overuse injuries i.e., an athlete may continue to train and compete but at lower intensity or efficiency. For this reason, sports medicine research has focussed on developing self-reported tools of which the best validated is the Oslo Sports Trauma Research Tool (OSTRC) [10]. When employing such tools, an interesting pattern emerges. Typically, 20-30% of athletes reported a negative impact of their oral health on training or performance. More specific questions reveal: 35% athletes reported difficulties eating or drinking, 17% difficulties smiling, laughing, or showing teeth without embarrassment and 15% difficulties relaxing. Therefore, elite and professional athletes commonly report a negative impact of their oral health on their training and performance and in addition, more specific effects on wellbeing.

Identifying the pathways by which oral health affects performance is ongoing research. Such pathways could include physical effects such as raised systemic inflammation resulting from oral disease as well as psychosocial effects through pain, self-reported bleeding, and appearance.

No direct evidence has yet been published that identifies an impact of oral inflammation on sport performance. However, in non-athlete populations, periodontitis has been found to be a risk factor

for low cardiorespiratory fitness in sedentary non-smoking men [11]. These findings were confirmed in a cross-sectional analysis of two large German cohorts, including 1,639 and 2,439 subjects, respectively [12]. Probing pocket depth (PPD) was consistently associated with VO<sub>2</sub>peak as well as exercise duration. Interestingly, this association remained significant after restricting the analysis to cardiorespiratory healthy participants only. Similarly, periodontitis was associated with poorer physical activity test results in male police officers following adjustment for age, BMI, and regular exercise [13]. There was a dose response with poorer outcomes associated with severity of periodontitis.

In addition to raised systemic inflammation another pathway for impact on performance is through psychosocial impacts. Many researches have published statistically significant associations between a number of the factors described earlier with negative performance impacts [4-6,12,13]. These factors are also commonly associated with impacts on oral health related quality of life and therefore highly plausible that they might affect performance. In relation to the other self-reported impacts mentioned earlier regarding common effects of poor oral health in athletes on eating and drinking, sleep and relaxation, there are additional possibilities that there could be negative effects on nutrition and sleep hygiene both of which are key to supporting athlete performance, recovery, and wellbeing [14].

## Conclusion

Oral health is closely linked to systemic health. For this reason, physical activity plays a crucial role in both general well-being and sports performance in professional athletes.

Performance impacts from oral health are hard to research because of their relatively lower magnitude although important within high performance sport. Nevertheless, negative performance impacts are common. The evidence is clearest for psychosocial effects arising from poor oral health. In non-athlete populations, periodontal inflammation is associated with reduced measures of fitness and may point to a further effect of the well-established phenomenon of raised systemic inflammation caused by periodontitis and other oral infections [5,6].

## Acknowledgement

None.

## Conflict of Interest

No conflict of interest.

## References

1. Needleman I, Ashley P, Petrie A, Fortune F, Turner W, et al. (2013) Oral health and impact on performance of athletes participating in the London 2012 Olympic Games: a cross-sectional study. *Br J Sports Med* 47(16): 1054-1058.
2. Locker D (1988) Measuring oral health: a conceptual framework. *Community Dent Health* 5(1): 3-18.
3. Cullinan MP, Seymour GJ (2013) Periodontal disease and systemic illness: will the evidence ever be enough? *Periodontol* 2000 62(1): 271-286.
4. Needleman I, Ashley P, Fine P, Haddad F, Loosemore M, et al. (2014) Consensus statement: Oral health and elite sport performance. *Br Dent J* 217(10): 587-590.
5. Needleman I, Ashley P, Meehan L, Petrie A, Weiler R, et al. (2016) Poor oral health including active caries in 187 UK professional male football players: clinical dental examination performed by dentists. *Br J Sports Med* 50(1): 41-44.
6. Needleman I, Ashley P, Weiler R, McNally S (2016) Oral health screening should be routine in professional football: a call to action for sports and exercise medicine (SEM) clinicians. *Br J Sports Med* 50(21): 1295-1296.
7. White DA, Tsakos G, Pitts NB, Fuller E, Douglas GV, et al. (2012) Adult Dental Health Survey 2009: common oral health conditions and their impact on the population. *Br Dent J* 213(11): 567-572.
8. Botelho J, Vicente F, Dias L, Júdice A, Pereira P, et al. (2021) Periodontal Health, Nutrition and Anthropometry in Professional Footballers: A Preliminary Study. *Nutrients* 13(6): 1792.
9. Ashley P, Di Iorio A, Cole E, Tanday A, Needleman I (2015) Oral health of elite athletes and association with performance: a systematic review. *Br J Sports Med* 49(1): 14-19.
10. Clarsen B, Myklebust G, Bahr R (2013) Development and validation of a new method for the registration of overuse injuries in sports injury epidemiology: the Oslo Sports Trauma Research Centre (OSTRC) overuse injury questionnaire. *Br J Sports Med* 47(8): 495-502.
11. Eberhard J, Stiesch M, Kerling A, Bara C, Eulert C, et al. (2014) Moderate and severe periodontitis are independent risk factors associated with low cardiorespiratory fitness in sedentary non-smoking men aged between 45 and 65 years. *J Clin Periodontol* 41(1): 31-37.
12. Gallagher J, Ashley P, Petrie A, Needleman I (2019) Oral health-related behaviours reported by elite and professional athletes. *Br Dent J* 227(4): 276-280.
13. Oliveira JA, Hoppe CB, Gomes MS, Grecca FS, Haas AN (2015) Periodontal disease as a risk indicator for poor physical fitness: a cross-sectional observational study. *J Periodontol* 86(1): 44-52.
14. Collins J, Maughan RJ, Gleeson M, Bilsborough J, Jeukendrup A, et al. (2021) UEFA expert group statement on nutrition in elite football. Current evidence to inform practical recommendations and guide future research. *Br J Sports Med* 55(8): 416.