

**Research Article**

Copyright © All rights are reserved by Saccomanni Bernardino

Outcomes for Tubs (Traumatic Unidirectional Bankart Surgery (Arthroscopic Treatment))

Saccomanni Bernardino**Orthopaedic and Trauma Surgery, Viale Regina Margherita, 70022, Altamura (Bari), Italy****Corresponding author:** Saccomanni Bernardino, Orthopaedic and Trauma Surgery, Viale Regina Margherita, 70022, Altamura (Bari), Italy**Received Date:** December 10, 2025**Published Date:** December 17, 2025**Abstract**

Background/Objectives: Despite the improvements in the methods of arthroscopic stabilization of anterior shoulder instability, a recurrence rate of as high as 30% is reported in the literature. In this context, we report the outcome of arthroscopic Bankart repair in anterior shoulder instability, with the use of bio-absorbable suture anchors for patients that were followed up for at least two years from the date of surgery. The arthroscopic method offers a less invasive technique of Bankart repair for traumatic anterior shoulder instability. We would like to report the 2-year clinical outcomes of bio-absorbable suture anchors used in traumatic anterior dislocations of the shoulder.

Methods: Data from 79 shoulders in 74 patients were collected over 4 years (2005–2009). Each patient was followed up over a period of 2 years. The patients underwent arthroscopic Bankart repair using bio-absorbable suture anchors for their anterior shoulder instability. These surgeries were performed at a single institution by a single surgeon over the time period. The patients were assessed with two different outcome measurement tools. The University of California at Los Angeles (UCLA) shoulder rating scale and the Simple Shoulder Test (SST) score. The scores were calculated before surgery and at the 2-year follow-up. The recurrence rates, range of motion as well post-operative function and return to sporting activities were evaluated.

Results: SST results from the 12 domains showed a significant improvement from a mean of 6.1 ± 3.1 to 11.1 ± 1.8 taken at the 2-year follow-up ($p < 0.0001$). Data from the UCLA scale showed a pre- and post-operative mean of 20.2 ± 5.0 and 32.4 ± 4.6 respectively ($p < 0.0001$). 34 had excellent post-operative scores, 35 had good scores, 1 had fair score and 3 had poor scores. 75% of the patients returned to sports while 7.6% developed a recurrence of shoulder dislocation or subluxation.

Conclusion: Arthroscopic Bankart repair with the use of suture anchors is a reliable treatment method, with good clinical outcomes, excellent post-operative shoulder motion and low recurrence rates.

Keywords: Outcomes; tubs; level of evidence: IV

Introduction

Recurrent shoulder dislocation or instability is common in young athletes. These injuries often occur during sports, preventing the individual from returning to these activities. The stability of

the glenohumeral joint is maintained by the glenoid labrum. This labrum creates a socket-deepening effect hence preventing any shoulder dislocations.

An avulsion of this anterior inferior labrum from the glenoid rim was first described by Perthes and Bankart in the early twentieth century [1,2]. Since then, several open and arthroscopic techniques have been described to address anterior shoulder instability. These procedures address both capsuloligamentous laxity and labral pathologies via a variety of instruments, suture passages, knot-tying techniques and fixation devices. With the debate continuing regarding the indications for arthroscopic shoulder stabilization, recent studies have shown favourable outcomes with regards to the arthroscopic method [3,4]. Moreover, with continuing criticisms with regards to the wide dissection, loss of external rotation, and post-operative pain associated with the open repair, the demand for arthroscopic surgery has increased over the last two decade.

However, despite advances in the understanding and techniques of arthroscopic surgery, failure rates have reported to be as high as 30%. As arthroscopic techniques have continued to evolve over the last decade, it is important to evaluate if these new techniques have resulted in an improved outcome. Our study aims to report and evaluate the pre-operative evaluation, thorough diagnostic arthroscopic examination for concomitant pathology, surgical techniques and the post-operative therapy programme for a successful outcome of arthroscopic Bankart repair with the use of bio-absorbable suture anchors for patients that were followed up for at least two years from the date of surgery.

Materials and Methods

From 2005 to 2009, a total of 79 shoulders in 74 patients underwent arthroscopic Bankart repair for recurrent anterior glenohumeral instability by a single surgeon at our institution. Five patients underwent bilateral shoulders stabilization. We conducted a retrospective analysis of a prospectively collected data after approval was sought for our study protocol from our hospital's ethics committee [5]. patients were lost to follow-up for UCLA analysis and [6] patients did not complete the SST questionnaire because at last the patients refused the questionnaire. Inclusion criteria for surgery included recurrent anterior glenohumeral subluxation or dislocation after an initial episode of traumatic anterior shoulder dislocation, a Bankart lesion confirmed by arthroscopic examination or ultrasound or Magnetic resonance imaging (MRI) and arthroscopic Bankart repair done using bio-absorbable suture anchors. The exclusion criteria were posterior instability, multidirectional instability, Hill-Sachs lesions more than 25% of the humeral head and bony Bankart lesion more than 25%. The degree of structural bony lesions was evaluated during arthroscopy, and patients demonstrating an engaging hill sac or an inverted pear glenoid were taken to have significant bony loss. All patients demonstrated a positive apprehension test as well as a load and shift test. All patients had pre-operative radiographs with an anterior-posterior, lateral, axillary and scapular-Y views taken. Magnetic resonance arthrograms were performed in patients with equivocal findings. The patients were included in the study after obtaining written, informed consent.

Two different outcome scoring measures were used to evaluate the effectiveness of the arthroscopic Bankart repair. The shoulder

rating scale of University of California Los Angeles (UCLA)6 and the simple shoulder test (SST) [7]. The SST consisted of a series of 12 yes-no questions, measuring pain and function of the shoulder through assessing the patient's ability to perform 12 simple tasks with the affected shoulder. The maximum total score was 12 points, with a higher score indicating better function. The UCLA was used to evaluate the patient's pain, function, forward flexion, strength and patient satisfaction. These five items are rated on ordinal scales of different lengths and scoring points. The maximum total score possible is 35, with a higher score indicating better shoulder function. We assigned a score of 34-35 points as excellent, 29-33 points as good, 21-28 as mild, and 20 or less as poor.

The UCLA and SST were chosen based on reproducibility, practicability, ease of use and ease of incorporation in clinical practice. We believe that they were the most responsive scoring systems and also most accurately reflect the outcomes of the surgery by assessing the tasks the patients are able to perform with the shoulder [8]. The UCLA has also shown to have a low inter-observer variability [9], while the SST has also been shown to satisfy the American Shoulder and Elbow Surgeons recommended attributes for a shoulder function assessment form [10]. Furthermore, these 2 outcome scores have also been used on numerous occasions in evaluating instability of the shoulder [11]. Data analysis comparing the pre-operative and post-operative UCLA scores were done using the Wilcoxon matched pairs test and data comparing the before and after surgery outcomes for the SST were done using the Unpaired t test. A value of $p < 0.001$ was taken as significant. All patients were followed up in clinic at 2 weeks, 1 month and then at 6 monthly intervals. All patients had a minimum of 2 years follow-up. Pre and post-operative range of motion, function and return to sports were recorded. Treatment failure was regarded as recurrent shoulder dislocation, any sensation of subluxation, or instability preventing return to full activity or requiring a further stabilizing procedure.

Surgical Procedure

All operations were performed with the use of a standardised technique by the same surgeon. After induction of a general anaesthesia, the patient was placed in a beach chair position and a thorough examination under anaesthesia was performed to assess the magnitude and direction of instability. The shoulder was prepared and draped in a sterile manner, and the bony landmarks were marked carefully to maintain orientation throughout the procedure. A standard posterior viewing portal was established approximately 2 cm inferior and 1 cm medial to the acromial angle. Two anterior portals were established using outside-in technique with a spinal needle to establish the most appropriate placement of the cannulas. The anterosuperior portal was made in the rotator interval just inferior to the anterior edge of the acromion, and the anterior mid-glenoid portal was made just over the superior border of the subscapularis tendon. A small cannula was inserted into the anterosuperior portal, and a large-diameter threaded cannula was placed in the anterior mid-glenoid portal. Complete diagnostic arthroscopy was done through the posterior and anterior portals, with assessment of the glenoid labrum, capsule, rotator cuff and

the humeral head for possible Hill-Sachs lesions. Rotator interval closure was not performed and any other tears of the glenoid labrum were repaired.

The Bankart lesion was mobilised from the anterior glenoid surface using a periosteal elevator. The goal was to mobilise the labrum such that it could be shifted superiorly and laterally. The glenoid neck was lightly abraded using a rasper. All suture anchors used were from obtained from Arthrex. The Bio-suture Tak is a 3 mm diameter by 13 mm long bio-absorbable "push-in" anchor with a moulded-in suture eyelet ideally suited for soft tissue attachment to bone in the shoulder joint where a small anchor profile with high pull-out strength is required. This suture anchor is moulded from PLDLA poly (l-lactide-co-D, l-lactide), a non-crystalline, bio-absorbable copolymer. Fig. 1 demonstrates the suture anchor used. The first anchor was placed at the 5.30 o'clock position, on the glenoid articular surface 3 mm from the articular edge. We believe this is essential in recreating the labral bumper, re-establishing the concavity-compression effect and also tensioning the inferior glenohumeral ligament. The most inferior placement would ideally be placed at the 6 o'clock position; however, this often is not possible

due to limitations in the placement angle. The suture anchor used requires drilling a pilot hole or using a punch to create the pilot hole prior to impaction of the implant to a countersunk position in the bone. A suture passer is then passed under the Bankart lesion. The suture strand of the suture anchor nearer the labrum was brought out through the anterosuperior portal, and in turn through the labrum in a retrograde fashion using the suture passer and retrieved from the mid-glenoid portal. This suture limb remained as the post during suture tying and this would ensure that the knot rest of the capsular side of the glenoid labrum and not on the articular side. This technique would effectively push the labrum up towards the glenoid socket, restoring labral height [12] and thereby recreating the labral bumper. Lazarus et al showed in a cadaveric study that by reducing the labral height by 80%, the resultant stability of the joint was decreased by 60% and that restoring of the labral height was paramount in restoring stability of the glenohumeral joint [13]. Hence our goal through the above techniques described through anatomical restoration of labral complex we hope to restoring tension in the anterior inferior glenohumeral ligament and achieve stability of the glenohumeral joint (Figure 1).



Figure 1: Demonstrates the suture anchor used.

The second and third suture anchors were done at the 4.30 and 3.30 o'clock positions in the same manner. The sutures were tied using the Tennessee slider knot, which is easy to tie, has a low-profile and possesses good holding strength [14]. When there was evidence of anteroinferior capsular laxity, the suture passer would be passed through the peri labral capsule one cm anterior and one cm inferior to the Bankart lesion to plicate the redundant capsule. This laxity is assessed by the ability to pass the arthroscope between the humeral head and the glenoid at the level of anterior band of the inferior glenohumeral ligament. This drive-through sign is considered to be diagnostic of shoulder laxity or instability [15].

Post-operatively, the patients were placed in a sling for six weeks. They were allowed to do pendular motion exercises for the

first three weeks, followed by elevating the elbow to shoulder level (forward active flexion to 90°) from the third to the sixth week. They were also taught to do isometric rotator cuff exercises during these six weeks. Full shoulder mobilisation was allowed after six weeks. Sport activities were allowed at three months and contact sports at four months (Table 1).

Table 1 demonstrates the biostatistics of the patients in this study. There were no complications with regards to the arthroscopic technique. No bleeding, infection, compartment syndrome or neurological compromise was observed post-operatively. The most common associated injury was a Hill-Sach's lesion. This occurs as the posterior aspect of the humeral head impacts against the anterior glenoid, when the shoulder is dislocated anteriorly.

Table 1: Biostatistics of the patients who underwent Arthroscopic Bankart repair.

Average Age (range) (years)	24.85 (13-44)
Gender	
Male	74
Female	1
Number of shoulders*	79 (5)
Mean number of dislocations before surgery (range)	11.17 (1-100)
Mean duration of operative time (range)	64.56 (35-145)
Mean pre-operative range of external rotation (range)	79.60 (60-90)
Mean post-operative range of external rotation (range)	81.39 (60-90)
Mean number of suture anchors (range)	2.87 (2-3)
Operative finding (Number of shoulders)	
Bankart lesion	79
Hill-Sachs lesion (mild grade)	10
Chondrolabral Lesion	1
Bony Bankart lesion	>25% 0
SLAP Lesion	2
Lax anteroinferior capsule (required capsular plication)	11

Results

The Simple Shoulder test (SST) showed a total of 73 responses out of the 79 shoulders that were operated on. The SST showed

statistically significant improvement ($p < 0.0001$) from the pre-operative scores from a mean and standard deviation (SD) of 6.06 ± 3.12 with a range from 0 to 8 to a mean and SD of 11.08 ± 1.78 and a range from 4 to 12 (Table 2).

Table 2: UCLA outcome scores in patients after an arthroscopic Bankart repair with suture anchors.

	Mean and SD before surgery (n = 73)	Mean and SD after surgery (n = 73)	p value (Unpaired T)
Pain	5.84 ± 2.33	8.79 ± 1.62	<0.0001
Function	5.74 ± 2.54	9.18 ± 1.69	<0.0001
Active Forward Flexion	4.44 ± 0.91	4.95 ± 0.23	<0.0001
Strength of forward flexion	4.05 ± 1.10	4.79 ± 0.55	<0.0001
Satisfaction of patient	0	4.73 ± 1.15	<0.0001
Total	20.21 ± 4.98	32.44 ± 4.60	<0.0001
Pre-operative	Post-operative		
Number of shoulders who scored poor	36	3	
Number of shoulders who scored fair	32	1	
Number of shoulders who scored good	5	35	
Number of shoulders who scored excellent	0	34	

Table 2 demonstrates the scores from the UCLA evaluated the patient's pain, function, active forward flexion, strength of forward flexion and satisfaction of the patient. Total UCLA score showed an improvement from a mean and SD of 20.21 ± 4.98 before surgery to 32.44 ± 4.60 post surgery, with 69 shoulders achieving excellent or good scores (94.5%), 1 having a fair score (1.5%), and 3 having poor scores (4.1%). All patients demonstrated good range of motion

with a mean and SD external rotation of $81.39 \pm 8.12^\circ$.

A total of 6 shoulders in 5 patients had a recurrence of shoulder instability. Of the 6, 4 of the recurrence of dislocation were due to sporting activities, while the causes of dislocation of 2 shoulders were unknown. 75% of the patients returned to previous sporting activities, while the remainder felt they could not return because

they were afraid of a recurrence. All of the patients apart from those who developed a recurrence demonstrated a negative load and shift as well as a negative anterior apprehension test on post-operative clinical examination. Patients were also asked to rate the feeling of stability of their shoulder pre- and post-operation on a scale of 0-10, with 10 being the most unstable. Mean shoulder instability score was 7.33 before surgery and 1.89 after surgery.

No correlation could be established between the age, gender, frequency of dislocation, duration from first dislocation to surgery and the rate of recurrence. Although Voos and his colleagues found associated ligamentous laxity and age under 25 to be risk factors for recurrence, these factors could not be established in our study [16].

Discussion

Historically, arthroscopic repair for the treatment of the Bankart lesion has been less satisfactory than the open technique. However, many of these arthroscopic techniques described were using trans glenoid sutures or bio-absorbable tacks [17]. In last few years, newer techniques involving suture anchor fixation and capsular plication have started to evolve, with promising results. These suture anchors have increasingly been using in labral repair and capsulolabral reconstruction [18]. Our study has shown that patients undergoing arthroscopic repair with these suture anchors have excellent clinical outcomes and similar recurrence rates as compared to open surgery.

Suture anchors are low-profile fixation devices that minimize articular surface damage of the humeral head, offering anatomic reconstruction of the glenoid labrum as well as the glenohumeral ligament complex. These suture anchors may be inserted either open or arthroscopically, with the aim of re-attaching the anterior inferior labrum along with the ligaments to the glenoid labrum. Knots are placed on the capsular side of the Bankart lesion, recreating the socket-deepening bumper effect of the labrum and hence restoring the concavity-compression mechanism of the glenoid labrum on the humeral head [19]. Any redundant or loose capsule is also addressed during the same operation, allowing one to address any capsular laxity, restoring tension in the anterior inferior glenohumeral ligament and stability to the glenohumeral joint.

The arthroscopic Bankart repair offers many advantages when compared to the open technique. It offers a minimally invasive approach with less surgical trauma and blood loss, with improvements in operating time, peri-operative morbidity, narcotic use, hospital stay, time loss from work and decrease number of complications together with a lower cost of surgery [20]. We have also shown that post-operative range of motion is not sacrificed for the sake of stability, with a mean and standard deviation of $81.39 \pm 8.12^\circ$ of external rotation. This allows the patients to return to sports or return to physically demanding jobs.

The introduction of bioabsorbable suture anchors also simplifies any revision surgery, reducing concerns regarding infected implants [21] and anchor migration leading to articular cartilage

damage [22]. During surgery, either two or three suture anchors are inserted, depending on the size of the Bankart lesion. Our results showed that patients who had only two suture anchors did not have a higher rate of recurrence. Patients with anteroinferior capsular laxity were treated accordingly by pinch tuck capsular plication as described earlier. Although some studies have shown that the presence of capsular laxity may affect the outcome of arthroscopic stabilization [23], while others have suggested that the elastic deformation of the glenohumeral ligament at the time of injury prevents the same degree of structural damage [24], we do not consider Bankart lesions associated with capsular laxity a contraindication to arthroscopic surgery. On the contrary, capsular plication can be done arthroscopically to address the issue of anteroinferior capsular laxity and this significantly augments the stability achieved with Bankart repair.

The majority of our patients were young physically active individuals, who engage in either vigorous sports or high demand jobs. Satisfactory range of motion, especially external rotation allows for performance during sports as well as proper functioning for activities during daily living. Several other studies published also reported a good range of motion after arthroscopic repair, often even better than repair with the open technique [25].

The recurrence rate in our study was 7.6%, which is similar to other published studies. Recurrence rates using the open technique ranged from 0 to 22% [26]. Warner et al initially published discouraging results with the arthroscopic techniques for contact sport athletes [27]; however, with modern arthroscopic techniques, extremely strong suture anchors and secure repair techniques allowing the patients to undergo extensive rehabilitation our study and other supporting studies have shown early return to competitive sporting activities [28,29].

Conclusions

Arthroscopic Bankart repair with the use of suture anchors is a reliable treatment method, with good clinical outcomes, excellent post-operative shoulder motion and low recurrence rates.

Conflict of interest

No benefits in any form have been received or will be received from a commercial party related directly or indirectly to the subject of this article.

Reference

1. Perthes G (1906) On surgical procedures for habitual shoulder dislocation. *Dtsch Z Chir* 56: 149-151.
2. Bankart ASB (1923) Recurrent or habitual dislocation of the shoulder. *Br Med J* 1: 1132-1133.
3. Sperling JW, Smith AM, Cofield RH (2007) Patient perceptions of open and arthroscopic shoulder surgery. *Arthroscopy* 23(4):361-366.
4. Fabbriani C, Milano C, Demontis A, Fadda S, Ziranu F, et al. (2004) Arthroscopic versus open treatment of Bankart lesion of the shoulder: a prospective randomized study. *Arthroscopy* 20(5): 456-462.
5. Lo Ian KY, Parten Peter M, Burkhart S, Stephen M (2004) The inverted pear glenoid: an indicator of significant glenoid bone loss. *Arthroscopy* 20(2): 169-174.

6. Ellman H, Hanker G, Bayer M (1986) Repair of rotator cuff. End-result study of factors influencing reconstruction. *J Bone Jt Surg Am* 68(8): 1136-1144.
7. Lippitt SB, Harryman DT, Masten FA (1993) A practical tool for evaluating function: the simple shoulder test. In (Eds.), Masten FA, Fu FH, Hawkins RJ, editors. *The Shoulder: A Balance of Mobility and Stability*. American Academy of Orthopaedic Surgeons Rosemont pp. 501-518.
8. Godfrey J, Hamman R, Lowenstein S, Briggs K, Kocher M (2007) Reliability, validity, and responsiveness of the simple shoulder test: psychometric properties by age and injury type. *J Shoulder Elbow Surg* 16(3): 260-267.
9. Lam JJ, Ip FK, Wu WC (2000) Shoulder assessment systems: a comparison of three different methods. *Hong Kong J Med Sports*.
10. Richards RR, An KN, Bigliani LU (1994) A standardized method for the assessment of shoulder function. *J Shoulder Elbow Surg* 3(6): 347-352.
11. Sisto DJ (2007) Revision of failed arthroscopic Bankart repair. *Am J Sports Med* 35(4): 537-541.
12. Slabaugh MA, Friel NA, Wang VM, Cole BJ (2010) Restoring the labral height for treatment of Bankart lesions: a comparison of suture anchor constructs. *Arthroscopy* 26(5): 587-591.
13. Lazarus MD, Sidles JA, Harryman DT, Matsen FA (1996) Effect of a chondral-labral defect on glenoid concavity and glenohumeral stability. A cadaveric model. *J Bone Jt Surg Am* 78(1): 94-102.
14. Baumgarten KM, Wright RW (2007) Ease of tying arthroscopic knots. *J Shoulder Elbow Surg* 16(4): 438-442.
15. McFarland EG, Neira CA, Gutierrez MI, A J Cosgarea, M Magee (2001) Clinical significance of the arthroscopic drive-through sign in shoulder surgery. *Arthroscopy* 17(1): 38-43.
16. Voos JE, Livermore RW, Feeley BT, David W Altchek, Riley J Williams, et al. (2010) Prospective evaluation of arthroscopic bankart repairs for anterior instability. *Am J Sports Med* 38(2): 302-307.
17. Freedman Kevin B, Smith Adam P, Romeo Anthony A, Brian J Cole, Bernard R Bach (2004) Open bankart repair versus arthroscopic repair with transglenoid sutures or Bioabsorbable Tacks for recurrent anterior instability of the shoulder. *Am J Sports Med* 32(6): 1520-1527.
18. Rudzki JR, Purcell Derek B, Wright Rick W (2004) Options for glenoid labral suture anchor fixation. *Operative Techniques in Sports Medicine* 12(4): 225-231.
19. Lippitt S, Matsen F (1993) Mechanisms of glenohumeral joint stability. *Clin Orthop Relat Res* 291: 20-28.
20. Wang Conrad, Ghalambor Navid, Zarins Bertram, Jon J P Warner (2005) Arthroscopic versus open Bankart repair: analysis of patient subjective outcome and cost. *Arthroscopy* 21(10): 1219-1222.
21. Ticker JB, Lippe RJ, Barkin DE, Carroll MP (1996) Infected suture anchors in the shoulder. *Arthroscopy* 12(5): 613-615.
22. Berg EE, Oglesby JW (1996) Loosening of a biodegradable shoulder staple. *J Shoulder Elbow Surg* 5(1): 76-78.
23. Neri BR, Tuckman DV, Bravman JT (2007) Arthroscopic revision of Bankart repair. *J Shoulder Elbow Surg* 16: 419-424.
24. Habermeyer P, Jung D, Ebert T (1998) Treatment strategy in first traumatic anterior dislocation of the shoulder. Plea for a multi-stage concept of preventive initial management. *Unfallchirurg* 101(5): 328-341.
25. Fabbriani C, Milano C, Demontis A, Salvatore Fadda, Fabio Ziranu, et al. (2004) Arthroscopic versus open treatment of Bankart lesion of the shoulder: a prospective randomized study. *Arthroscopy* 20(5): 456-462.
26. Cole BJ, L'Insalata J, Irrgang J, Warner JJ (2000) Comparison of arthroscopic and open anterior shoulder stabilization: a two to six-year follow-up study. *J Bone Jt Surg Am* 82(8): 1108-1114.
27. Warner JJ, Goitz RJ, Irrgang JJ, Groff YJ (1997) Arthroscopic assisted rotator cuff repair: patient selection and treatment outcome. *J Shoulder Elbow Surg* 6(5): 463-472.
28. Tambe Amol, Badge Ravi, Funk Lennard (2009) Arthroscopic rotator cuff repair in elite rugby players. *Int J Shoulder Surg* 3(1): 8-12.
29. Flurin PH, Guillemette C, Guillo S (2007) Traumatic rotator cuff tears in rugby players. *J Traumatol Sport* 24: 203-206.