



Research Article

Copyright © All rights are reserved by Hatim MY Hamadnalla

Biochemical Properties and Nutritional Value of Balanites Aegyptiaca (Laloub) Seed Oil

Amna Elsiddig Ali¹, Babiker Mohammed², Hisham Abd Almonem Mahmoud Ali³, Inass Omer Mohamed Malik⁴, Mahdi Abd elmageed Mohammed Ali⁵ and Hatim MY Hamadnalla^{6*}

^{1,2,3,4}Department of Food Technology, University of Bahri, Sudan.

⁵Department of Biology and Technology, University of Bahri, Sudan

***Corresponding author:** Hatim M Y Hamadnalla Department of Biochemistry, College of Applied and Industrial Sciences, University of Bahri, Sudan.

Received Date: September 25, 2021

Published Date: October 07, 2021

Abstract

The main goal of this research was to study the physical and chemical properties of Balanites. Aegyptiaca (laloub) oil. Two kilograms of Balanites aegyptiaca fruit were purchased from local market in Omdurman, Sudan. Fruits were then crushed using a steel hammer and seeds were then obtained. One kilogram of seeds was collected and was then ground using grinding machine ground seeds was then eventually ready for further analysis. The seeds of Balanites aegyptiaca were collected, washed and prepared. The seed oil was extracted using soxhlet apparatus with n-hexane as solvent. The percentage yield of oil extraction was (41.9 %). The results revealed that B. aegyptiaca seed contain considerable percentage of moisture (3.27%), fat (41.9), protein (30.9%), fiber (11.34%), ash (3.55%) and carbohydrate (8.9%). Physicochemical analysis of the seed oil was conducted using standard procedures and the following results were obtained; density (0.9091 g/cm³), viscosity (42.71 %), refractive index (1.4734), iodine value (98.09 g/100g), saponification value (194.33 mg KOH/g) and peroxide value (3.08meq/kg), acid value (0.53 mg KOH/g), free fatty acid value (0.26 %) and un-saponification (3.08). This reveals that Balanites aegyptiaca seed oil could be a rich source of oil for domestic and industrial purposes if richly exploited.

Introduction

Balanites aegyptiaca is used for firewood, charcoal, poles, timber, utensils, tool handle, food, fodder, mulch, shade, wind break and gum. The plant may be grown for its fiber, oil and for medicinal values; it is also used in treatment of several diseases and disorders since ages. Balanites aegyptiaca seed kernel oil falls in the group of such oils termed as vegetable oils since it is extracted from a plant source. The chief importance of vegetable oils lies in their food value.

The seed kernel is considered as an extremely useful edible product, it is rich in oil, protein, minerals and it has been reported to be used for over thousands of years. (Vonmaydell, H.J. (1986)) [1] Oils are heterogeneous biochemical substances which have in common, the property of being soluble in most organic polar

solvents (chloroform, benzene, diethyl ether, etc.) and insoluble in water. The term lipid is the scientific name for fatty acids and similar chemicals often found in oils produced by living things. They may contain diethyl-glycerol, free fatty acids, phosphatides, sterols and fat-soluble vitamins like tocopherol, pigments, waxes and fatty alcohol are present but as minor components [2].

Oils are heterogeneous biochemical substances which have in common, the property of being soluble in most organic polar solvents (chloroform, benzene, diethyl ether, etc.) and insoluble in water. The term lipid is the scientific name for fatty acids and similar chemicals often found in oils produced by living things. They may contain diethyl-glycerol, free fatty acids, phosphatides, sterols and fat-soluble vitamins like tocopherol, pigments, waxes

and fatty alcohol are present but as minor components. Oils may be sourced from animal, vegetables or petrochemical. All oils and fats are made up of a mixture of these triglycerides

Soaps are compounds that consist of long chain of hydrocarbons attached with a carboxylic acid which is ironically bonded to the metal ion usually sodium or potassium, it is a combination of animal fat or plant oil with caustic soda. Soaps are therefore called the salts of fatty acids which are usually used as surfactants for washing, bathing and cleaning. Soaps are obtained by the treatment of vegetable oils or animal fats with a strong alkaline solution. The three molecules of fatty acids in the triglyceride gets attached to a single molecule of glycerol and results in a chemical reaction termed saponification.

The extracted oil is used for many uses and it is used in Western Sudan remaining cake is used as animal feed. Both fruits and seed were widely used in many countries during the dry season and drought periods including Nigeria (Lockett et al., 2000), Ethiopia, (Guinand and Lemessa, 2001) [3] and Sudan (Grosskinsky and Gullick, 2001). The need for vegetable oil is rising worldwide so it has to be to look for good sources for the production of high-quality oil can be exploited for industrial purposes This study was conducted on the physical and chemical properties of *Balanites aegyptiaca* (Laloub) seed oil as a good source for the production of oil. Species *Balanites aegyptiaca* (L.) Delile – desert

date (Saed A. Al-Thobaiti and Isam M. Abu Zeid, 2018) [4].

Material

Two kilograms of *Balanites aegyptiaca* fruit were purchased from local market in Omdurman, Sudan. Fruits were then crushed using a steel hammer and seeds were then obtained. One kilogram of seeds was collected and was then ground using grinding machine ground seeds was then eventually ready for further analysis.

Results and Discussion

Chemical composition of *B. aegyptiaca* seed

Table (1) shows the chemical composition of *Balanites aegyptiaca* seed. Moisture, fat, protein, fiber, ash and total carbohydrate contents were found to be 3.27, 41.9, 30.9, 11.34, 3.55, and 8.9% respectively moisture content was found to be 3.27% which is similar with that reported by Mohammed .s. (2005) [5]. fat content was noticed to be 3.10%.but it was higher than reported Mohammed .S. Protein content was found to be 30.9% which is lower than the 31.09% reported by Mohammed S. (2005). Fiber content was 11.34% which is lower than the 12.64% reported by Mohammed S. (2005) [5]. Ash content was noticed to be 3.55% which is higher than the 3.19% reported by Mohammed S. (2005) [5]. Total carbohydrate content was recorded to be 8.9 % which is higher than the 3.05% reported by Mohammed S. (2005) [5] (Table 1).

Table 1: Chemical composition of *B. aegyptiaca* seed.

Experiment	First	Second	Three	Mean
Moisture	3.18	3.27	3.36	3.27
Fat	41.57	42.3	41.97	41.9
Protein	30.67	31.08	31.18	30.9
Fiber	10.67	11.38	11.97	11.34
Ash	3.67	3.56	3.42	3.55
Total carbohydrate	10.41	8.41	8.1	8.88

Physical properties of *B. aegyptiaca* oil

Table (2) shows that the physical properties of *Balanites aegyptiaca* oil. The viscosity, density, refractive index and color were found to be 42.71, 0.90913, 1.4734 and (red 5.1, yellow 35.5, blue 0.00) respectively.

Viscosity was recorded to be 42.71 which were higher than the 19.63 and 34cpreported by Mohammed S. (2005) [5], Babagana et.al (2010), respectively. Density was noticed to be 0.9091 which its similar to 0.9109 reported by Mohammed S. (2005) [5]. Refractive index was found to be 1.473 which is similar to 1.483 that reported by Mohammed S. (2005) [5] (Table 2).

Table 2: Physical properties of *B. aegyptiaca* oil.

Experiment	First	Second	Third	Mean
Viscosity	42.67	42.82	42.67	42.71
Density	0.9016	0.9131	0.9127	0.9091
Refractive Index	1.4786	1.4751	1.4667	1.4734
Color Red	5.2	5.1	5.1	5.1
Color Yellow	35.8	35.6	35.3	35.5
Color Blue	0	0	0	0

Chemical properties of *B.aegyptiaca* oil:

Table (3) shows that the chemical properties of *Balanitesaegyptiaca* oil the free fatty acid, peroxide value, saponification, iodine value, un-saponification, and acid value were found to be 0.26, 3.08, 194.33, 98.09, 3.08, and 0.53 respectively. Free fatty acid was found to be 0.26% which was lower than the 2.8 and 1.84% reported by both Babagana et.al (2011) [6] and Manj (2013) [7] respectively. Peroxide value was noticed to be 3.08(mgKO) which lower than the 6.0 and 8.0 reported by Manji(2013) [7] and Babeker (2013) [8] respectively. Saponification was reported to be 194.33 (mgKOH/g) which was higher than the 168.80, 174.5, 168.3 and 182.80 (mgKOH/g) reported by Manji (2013), Babagana gutti (2011) [6], Babeker (2013) [8] and Okia (2013) respectively. Iodine was found to be 98.09 mg I₂/g which higher than the 76.8, 56.4 and lower than 98.28 mg I₂/g reported by Manji (2013) [7], Babaganagutti (2011) [6-15] and Okia (2013) respectively. Acid value was noticed to be 0.53(mgKOH/g) which is similar to reported by Okia (2013) but its lower than the 2.08 (mgKOH/g) reported by Babeker [15-22]. Un-saponification was found to be 3.08 (Table 3) [23-29].

Table 3: Chemical properties of *B. aegyptiaca*oil.

Third	Second	First	Mean	Experiment
0.27	0.24	0.29	0.26	FFA
0.55	0.48	0.58	0.53	Acid value
3.26	2.88	3.11	3.08	Peroxide value
193.67	195.81	193.51	194.33	Saponification value
3.27	2.86	3.11	3.08	Un-saponification value
98.41	98.36	97.51	98.09	Iodine value

Conclusion

The results obtained from the preliminary investigation carried out in this work revealed that *B.aegyptiaca* seed oil is an economically viable oil source because its oil content was found to be high. Also, the oil parameters showed that the oil was composed of moderately long chain fatty acids with a degree of unsaturation, making it a good feedstock for domestic and industrial purposes [23-29].

Acknowledgement

None.

Conflict of Interest

No conflict of interest.

References

- Vonmaydell HJ (1986) Trees and Shrubs of the Sahel: Their Characteristics and Uses. Eschborn, GTZ, Germany: 525.
- Zang CU, Jock AA, Garba HI, Chindo YI (2018) Application of Desert Date (*Balanitesaegyptiaca*) Seed Oil as Potential Raw Material in the Formulation of Soap and Lotion. American Journal of Analytical Chemistry 9: 423-437.
- Guinand Y, lemessa D (2001) Wild Food Plants in Ethiopia: Reflections on the Role of Wild Foods and Famine Foods at a Time of Drought. In:

Kenyatta C, Henderson A (Eds.), the Potential of Indigenous Wild Foods. Workshop Proceedings, USAID/OFDA, Mombassa, Kenya

- Saed A Al-Thobaiti, Isam M Abu Zeid (2018) Medicinal Properties of Desert Date Plants(*Balanitesaegyptiaca*) – An Overview Department of Biological Sciences, Faculty of Sciences, King Abdulaziz University, P.O. Box 139109, Jeddah 21323, Saudi Arabia.
- Mohammed S Elmustafa (2015) Fregon Physicochemical Properties of *Balanitesaegyptiaca* (Laloub) Seed Oil November.
- Babagana Gutti, Shittu S Bamidele, Idris M Bugaje (2011) characterization and Composition of *Balanitesaegyptiaca* seed oil and its potential as biodiesel feed stock in Nigeria. Journal of Applied Phytotechnology in environmental sanitation1(1): 29-35.
- Manji AJ, Sarah EE, Modibbo UU (2013) Studies on the potentials of *Balanitesaegyptiaca* seed oil as raw material for the production of liquid cleansing agents. Department of Industrial Chemistry, Modibbo Adama University of Technology, P. M. B. 2076, Yola, Adamawa State, Nigeria.
- Babeker MA (2013) Physicochemical Properties of Laloub Seed oil. M. Sc. 29-45. Faculty of Agriculture, University of Khartoum.
- Ibrahim AM (2016) Extraction Oils from seeds of Baobab (*Ophelussitularius*) and *Batanites* (*Aegilopseniculata*)
- <https://en.m.wikipedia.org/wiki/batanites-aegyptiaca>. Genus: *Batanites* Delile U.S. National plant Germplasm system.(2016).
- Richard K, Francis O (2017) *Batanites Aegyptiaca* multi-purpose tree WWW.southworld.net/batanites-aegyptiaca-a-multi-purpose-tree.
- Fadi KM (2013) *Balanitesaegyptiaca*(L.) Del.: A multipurpose Fruit Tree on the Savanna Zone of Western Sudan Agricultural Research Corporation, El-Obied Research Station, P.O. Box 429, El-Obied, Sudan.
- Anon YM (2004) Final report, South Kordofan range management strategy study: 38. Yam Consultancy and Development. Co. Ltd. Sudan, Khartoum.
- Lazim AM (2007) Proximate composition of some fodder trees in the South Kordofan State. Ministry of Science and Technology Agricultural Research Corporation, Proceedings the 41 and 42 meetings of the national crop husbandry committee, Wad Medani.
- Doughari JH, Pukuma MS, De N (2007) Antibacterial effects of *Balanitesaegyptiaca*(L) Dell and *Moringaoleifera*Lam on *Salmonella typhi*. African Journal of Biotechnology 6(19): 2212-2215.
- Mordechay H, Zeev W, Sarina G (2008) Production of Biodiesel from *Balanitesaegyptiaca*. Browdy and Neimarry, P.L.L.C; 624, Washington, Dc, US.
- Garba HI, Chindo YI (2018) Application of Desert Date (*Balanitesaegyptiaca*) Seed Oil as Potential Raw Material in the Formulation of Soap and Lotion. American Journal of Analytical Chemistry 9: 423-437.
- Adebayo SE, Orhevba BA, Adeoye PA, Fase OJ, Musa JJ (2012) Solvent Extraction and Characterization of oil from African Star Apple (*Chrysophyllum albidum*) Seeds. International Journal of Academic research 3: 178-183.
- Elseed AMAF, Amin AE, Khadija AA, Ali JS, Hishinum M, et al. (2002) Nutritive Evaluation of Some Fielder Tree Species during the Dry Season in Central Sudan. Asian Aust. Journal Animal Science 15: 844-850.
- David P (1970) the chemical analysis of foods.
- <http://www.bangor.ac.uk/~afs032/Fruittrees.html>lit.
- Pereira s (2014) Freitas on 13 November 2014 modified Method for the determination of Unsaponifiable Matter in Oils and Fats.
- Author CC (1995) Lipid-Based Fats Substitutes. Critical Review Science Nutrients Journal 35: 4-5.
- Feranil AB, Dauzo PL, Kuzawa CW, Adair LS (2011) Coconut Oil Predicts a Beneficial Lipid Profile in Pre-Menopausal Women in the Philippines. Asian pacific Journal of clinical Nutrition 20: 190-195.
- Miller S (1994) The Natural Soap Book. Storey Publishing, North Adams: 45-50.
- AOAC (2008) Official Methods of Analysis, Association of Official Analytical Chemists, Washington D.C. (15th ed).

27. Pearson D (1981) *The Chemical Analysis of Food* (8Ed) J. A. Churchill London, pp. 535.
28. Aremu MO, Olonisakin A, Bako DA, Madu PC (2006) Compositional studies and physicochemical characteristics of Cashew nut flour. *Pak J Nutr* 5: 328-333.
29. Chineye GC, Akugwo EJ, Chineye NI, Ugbogu AE (2009) Nutritional value of *Balaniteaegyptiacaseed* (wild bottle gour) from South- Eastern Nigeria. *Parkistan Journal of Nutrition* 8 (3): 284-287.