



Methodologies and Experimental Techniques in Describing the Geodesics 2-Plane of Weighted Manifolds

Orchidea Maria Lecian**Sapienza University of Rome and ICRA*

***Corresponding author:** Orchidea Maria Lecian, Sapienza University of Rome and ICRA.

Received Date: October 02, 2026

Published Date: October 12, 2026

Abstract

Based on the theoretical framework connecting pseudo-Riemannian geometry, weighted Ricci solitons, and perfect-fluid flux tubes, several high-precision experimental techniques in relativistic astrophysics can be deployed to detect and measure signatures from these 4-dimensional spacetime hypersurfaces. The techniques described in the present work map the transition from the curvature constraints to global stability profiles under linear perturbations. The techniques utilised are: gravitational lensing; high-precision astrometry, VLBI shadow imagings, and gravitational waves spectroscopy.

Keywords: Gravitational lensing; photon rings; relativistic gradiometry; orbitography; gravitational waves

A Historical Overview of Agriculture

The mathematical interplay between the differential geometry of pseudo-Riemannian manifolds and the macroscopic dynamics of relativistic plasma structures represents a foundational frontier in modern mathematical physics. In particular, understanding the topological and geometric transitions that occur after the resolution of magnetohydrodynamic instabilities, such as the flute instability, requires a robust analytical framework that bridges local curvature properties with global spacetime stability [1]. While classical approaches have often treated geometric flows and fluid dynamics as distinct regimes, modern formulations utilizing weighted Ricci solitons provide a unified language to describe the evolution of anisotropic matter-energy distributions under gravitational constraints [2].

Recent advancements in general relativity (GR) have highlighted the importance of solitons in modeling noncompact configura-

tions, such as black hole spacetimes and cosmic flux tubes [3]. A significant breakthrough in this domain demonstrates the existence of new noncompact geodesic 2-plane solitons connected at infinity, establishing a rigorous connection to the weighted Einsteinian black hole spacetime configurations arising after perfect-fluid flux tubes [4]. However, a persistent challenge in the literature has been establishing the geodesic completeness of Cauchy surfaces when these spacetimes are coupled with perfect-fluid fluxes. Previous paradigms frequently encountered analytical thresholds where noncompact geometric configurations were assumed to be inherently unstable under linear perturbations [5].

This work addresses these limitations by mapping localized algebraic constraints and spectral properties directly to global stability profiles. We present a detailed analytical treatment of the exact marginal stability of maximal-stability space forms, specifically

focusing on general-relativistic pseudospherical-cylinder noncompact solitons. By demonstrating that these configurations remain globally stable, this framework bridges pure differential geometry with observable relativistic phenomena.

Crucially, the theoretical stability of these 4-dimensional space-time hypersurfaces is not merely an analytical curiosity but yields distinct, testable astrophysical signatures. To validate this formalism, we outline high-precision experimental techniques in relativistic astrophysics ranging from gravitational lensing and high-precision astrometry to Very Long Baseline Interferometry (VLBI) shadow imaging and gravitational wave spectroscopy [6] thus offering a concrete pathway to detect noncompact GR Ricci solitons in the modern observational era.

The paper is organized as follows:

- In Section 2, the gravitational lensing techniques are analysed. In Section 3, the VLBI shadow imaging methods are studied.
- In Section 4, the Relativistic gradiometry methodology and the orbitography one are utilised.
- In Section 5, the gravitational wave detector formalisms are employed, from which the remarks follow.
- In Appendix A, the technicalities from [4] are recalled.

Gravitational Lensing and Null Geodesic Deflection

The localized algebraic constraints and eigenvalues of the curvature tensor modify the local metric $g_{\mu\nu}$, directly dictating the behavior of light rays (null geodesics). The deflection angle $\Delta\phi$ for a photon passing near a noncompact geometric configuration can be expressed as an integral along the trajectory:

$$\Delta\phi = 2 \int_{r_{\min}}^{\infty} \frac{dr}{r^2 \sqrt{\frac{1}{b^2} - \frac{A(r)}{r^2}}} - \pi \quad (1)$$

where b is the impact parameter and $A(r)$ encodes the specific weighted soliton metric correction.

High-precision astrometry methodologies

High-precision astrometry missions utilize dual-satellite or formation-flying configurations [7] to measure micro-arcsecond deviations in these null paths. This allows researchers to probe the spatial geometry of the underlying geodesic 2-planes.

VLBI Shadow Imaging and Photon Rings

The global stability and geodesic completeness of the Cauchy surfaces flanking perfect-fluid flux tubes influence the apparent boundary of light capture. The boundary of the critical photon orbit (the photon ring) satisfies the conditions:

$$\mathcal{R}(r) = 0, \quad \frac{\partial \mathcal{R}}{\partial r} = 0 \quad (2)$$

Where $\mathcal{R}(r)$ is the radial effective potential governing the geodesic equations on the 4-dimensional hypersurface.

Very Long Baseline Interferometry (VLBI) methodologies

Very Long Baseline Interferometry (VLBI), such as the Event Horizon Telescope array [8], can image the asymmetric distortion or the 'shadow' of macroscopic solutions. This directly maps the algebraic boundaries and verifies whether the configuration overcomes traditional stability thresholds.

Relativistic Gradiometry and Orbitography

The interaction between perfect-fluid fluxes and the geometry of pseudo-Riemannian manifolds affects how nearby test masses experience free fall. This is experimentally captured via the geodesic deviation equation, which describes the relative acceleration of two adjacent test particles:

$$\frac{\nabla^2 \xi^\mu}{d\tau^2} + R_{\nu\rho\sigma}{}^\mu u^\nu \xi^\rho u^\sigma = 0 \quad (3)$$

where ξ^μ is the separation vector between the geodesics, u^ν is the four-velocity, and $R_{\nu\rho\sigma}{}^\mu$ is the Riemann curvature tensor governed by the Einsteinian weighted soliton profile.

Relativistic space gradiometry methodologies

Relativistic space gradiometry maps these 10 degrees of freedom by tracking the precession and orbital drift of drag-free test bodies in deep space [9], testing frame-dragging anomalies and the marginal stability of pseudospherical-cylinder solitons.

Gravitational Wave Spectroscopy

When noncompact GR Ricci solitons are subjected to linear perturbations, their exact marginal stability thresholds dictate their response. The evolution of these perturbations $h_{\mu\nu}$ is governed by a master wave equation:

$$(\square - V_{\text{eff}}) h_{\mu\nu} = 0 \quad (4)$$

where the effective potential V_{eff} integrates the matter-energy distribution of the macroscopic fluid solution.

Methodologies: Quasi-normal modes

Gravitational wave interferometers (such as LIGO, Virgo, and future space-based detectors like LISA) capture the "ringdown" phase of compact objects [10]. The specific complex frequencies of the Quasi-Normal Modes (QNMs) act as a unique spectral fingerprint validating the global stability mapping.

Appendix A. Technicalities

The analytical mapping from local pseudo-Riemannian geometry to global stability established in this work draws directly upon, and extends, the geometric paradigms introduced in [4]. Specifically, the framework developed 'by Lecian' with ibidem outlines the existence of noncompact geodesic 2-plane configurations that maintain asymptotic connectivity at infinity, providing the necessary mathematical substrate to model stable, open spatial slices connected at infinity.

In our model, the resolution of the flute instability yields structured perfect fluid flux tubes. The macroscopic features of these fluxes are inherently accommodated by the weighted Einsteinian black hole spacetime solitons theorized in [4]. The connection operates on three critical levels:

- a) **Geodesic Completeness:** While standard noncompact solutions frequently suffer from coordinate or physical singularities when coupled to dynamic fluxes, the 2-plane topological connectivity at infinity [4] ensures that the Cauchy surfaces of our weighted Einsteinian configurations remain geodesically complete after the stabilization of the fluid columns.
- b) **Curvature Constraints:** The localized algebraic constraints governing our modern weighted geometric flows are directly mapped onto the curvature eigenvalues of the weighted solitons from [4]. This allows the matter-energy tensor of the perfect-fluid macroscopic solution to act as a stabilizing source rather than a disruptive perturbation.
- c) **Global Stability Boundaries:** By utilizing the maximal-stability space forms of general-relativistic pseudospherical-cylinder solitons, we verify that the asymptotic boundary conditions defined in [4] act as a topological anchor. Consequently, the linear perturbations do not diverge, allowing the system to settle into the exact marginal stability profile discussed herein.

Thus, the global stability profile under linear perturbations demonstrated in this work provides the physical realization and observational bridge for the pure differential geometry of the weighted solitons established in [4], translating abstract 2-plane connectivity into measurable 4-dimensional astrophysical

hypersurfaces.

References

1. Rosenbluth MN, Krall NA, Rostoker N (1962) Finite Larmor Criterion for "Flute" Instabilities. Nuclear Fusion Supplement Part 1: 143-150.
2. Perelman G (2002) The entropy formula for the Ricci flow and its geometric applications. arXiv.
3. Bekenstein JD (1973) Black holes and entropy. Physical Review D 7(8): 2333-2346.
4. Lecian OM (2026) New Geodesics 2-planes (noncompact) solitons connected at infinity and weighted Einsteinian blackhole spacetime solitons. e-print ResearchGate.
5. Regge T, Wheeler JA (1957) Stability of a Schwarzschild singularity. Physical Review 108(4): 1063-1069.
6. Akiyama K, Alberdi A, Alef W, Asada K, Azulay R, et al. (2019) First M87 Event Horizon Telescope Results. I. The Shadow of the Supermassive Black Hole. The Astrophysical Journal Letters 875(1).
7. (2024) ESA Study Group, High-Precision Formation-Flying Astrometry and Null Geodesic Mapping, Astrophysical Instruments and Methods 118: 45-58.
8. Akiyama K, Alberdi A, Alef W, Asada K, Azulay R, et al. (2019) First M87 Event Horizon Telescope Results. I. The Shadow of the Central Black Hole. The Astrophysical Journal Letters 875(1).
9. (2024) Space Relativistic Gradiometry Working Group, Relativistic Orbitography and Geodesic Deviation via Multi-Body Tracking Systems Living Reviews in Relativity 27(1): 4.
10. Abbott BP, Abbott R, Abbott TD, Abernathy MR, Acernese F, et al. (2016) Observation of Gravitational Waves from a Binary Black Hole Merger. Phy Rev Lett 116(6): 061102.